

# **Iba1 Recombinant antibody**

Cat: B09001R Company: Haokebio

Uniprot ID:P55008 Applications: IHC:1:500-1:4000

Organism: Rabbit IHC-Polymer: 1:2000-1:16000

IHC-TSA:1:2500-1:20000

IF:1:50-1:200

WB:1:500-1:1000

# Species reactivity: Human Mouse Rat

Molecular Weight Calculation: 17 kDa Observed Molecular Weight: 17 kDa

# Background:

Ibal is a protein that exists in the cytoplasm, and it is highly evolutionarily conserved. It is also p ossibly identical to three other proteins, Iba-2, M RF-1 and daintain. However complete functional profiles of all three proteins and how they overla p is unknown. Iba1 is a 17-kDa EF hand protein t hat is specifically expressed in macrophages / mi croglia and is upregulated during the activation o f these cells. Iba1 expression is up-regulated in microglia following nerve injury, central nervous system ischemia, and several other brain disease s. In recent years, the possibility of a role for Iba 1 in cancer development has also been considere d. Significantly higher levels of Iba1 expression were found in hepatocarcinoma cell lines and in t issue compared to healthy samples. It has also be en shown that Iba1 expression can contribute to progression of cancer by inhibition of apoptosis i n cells.

# Protein full name:

allograft inflammatory factor 1

#### Synonyms:

AIF1, AIF 1, AIF-1, allograft inflammatory facto r 1, G1

### Immunogen:

Recombinant protein

# Isotype:

IgG

#### Subcellular location:

Cytoplasm, Cell projection

# Purity:

Affinity purification

#### Form:

Liquid

# Storage Buffer:

PBS with 0.02%sodium azide,100  $\mu$ g/ml BSA and 50% glyce rol.


## Storage:

Store at -20 °C for one year.

# **Experimental procedure:**


Antigen retrieval: Citrate buffer (pH 9.0) , Medium high heat for 8 minutes, stop for 7 minutes, medium high heat for 8 minutes. Incubate antibody,  $4^{\circ}\text{C}$  overnight. Secondary antibody: P oly-HRP Goat Anti-Rabbit & Mouse Universal Secondary Antibody, RT, 1h.

# Images:



Sample: Mouse brain, 4% PFA 12-24h.





Dilution of 1:1000 incubated at room temperatur e for 1.5 hours.

# Source of Reagents:

发表[中文论文]请标注:Iba1(B09001R)由杭州 浩克生物技术有限公司提供;

发表[英文论文]请标注:Iba1(B09001R) were kin dly provided by Hangzhou Haoke Biotechnolog y Co., Ltd.