

VDAC1/Porin Recombinant antibody

Cat: B31010T

Company: HaoKebio

Uniprot ID: P21796

Applications: IHC:1:500-1:2000

Organism: Rabbit

IHC-Polymer:1:2000-1:8000

Species reactivity: Human Mouse Rat

IHC-TSA:1:2500-1:10000

Molecular Weight Calculation: 31 kDa

IF:1:50-1:100

Observed Molecular Weight: 31 kDa

WB:1:2000-1:10000

Background:

VDAC1, also named as VDAC, porin 31HM, porin 31HL and plasmalemmal porin, belongs to the eukaryotic mitochondrial porin family. It adopts an open conformation at low or zero membrane potential and a closed conformation at potentials above 30-40 mV, to form a channel through the mitochondrial outer membrane and also the plasma membrane. Unlike other membrane transport proteins, porins are large enough to allow passive diffusion. Studies have shown that VDAC1 is subject to both phosphorylation and acetylation. The apparent molecular weight of VDAC1 is 30-37 kDa. Hypoxic conditions were found to trigger cleavage of the VDAC1 C-terminal to yield a 26-kDa truncated but active form. This antibody is specific to VDAC1.

Protein full name:

voltage-dependent anion channel 1

Synonyms:

Porin, VDAC1, Porin 31HL, PORIN 31 HL, Plasmalemmal porin

Immunogen:

Recombinant protein

Isotype:

IgG

Subcellular location:

Mitochondrion

Purity:

Affinity purification

Form:

Liquid

Storage Buffer:

PBS with 0.02% sodium azide, 100 µg/ml BSA and 50% glycerol.


Storage:

Store at -20 °C for one year.

Experimental procedure:

Antigen retrieval: Citrate buffer (pH 6.0), Medium high heat for 8 minutes, stop for 7 minutes, medium high heat for 8 minutes. Incubate antibody, 4°C overnight. Secondary antibody: Poly-HRP Goat Anti-Rabbit & Mouse Universal Secondary Antibody, RT, 1h.

Images:



Dilution of 1:10000 incubated at room temperature for 1.5 hours.

Source of Reagents:

发表[中文论文]请标注:VDAC1/Porin(B31010T)由杭州浩克生物技术有限公司提供;

发表[英文论文]请标注:VDAC1/Porin(B31010T) were kindly provided by Hangzhou Haoke Biotechnology Co., Ltd.