

# **GAPDH Recombinant antibody**

Cat: B31001T Company: HaoKebio

**Uniprot ID:**P04406 **Applications:** IF/ICC: 1:500-1:2000

**Organism:**Rabbit WB:1:5000-1:50000

Species reactivity: Human Mouse Rat

Molecular Weight Calculation: 36 kDa

Observed Molecular Weight: 36 kDa

## Background:

Glyceraldehyde-3-phosphate dehydrogenase (GA PDH) catalyzes the phosphorylation of glycerald ehyde-3-phosphate during glycolysis. GAPDH p articipates in nuclear events including transcripti on, binding RNA, RNA transportation, DNA repl ication, DNA repair and apoptosis. Being stably and constitutively expressed at high levels in mo st tissues and cells, GAPDH is considered a hous ekeeping protein. It is widely used as a control fo r RT-PCR and also loading control in electrophor esis and Western blotting. GAPDH is normally e xpressed in cellular cytoplasm or membrane, but can occasionally translocate to the nucleus after t he addition of post-translational modifications su ch as S-nitrosylation. This antibody is raised agai nst full length GAPDH of human origin. It can re cognize the 36 kDa GAPDH protein in most cells /tissues. In addition, a band below 36 kDa can al ways be detected as the isoform or spliced produ ct of GAPDH. Please note that some physiologic al factors, such as hypoxia and diabetes, increase GAPDH expression in certain cell types.

## Protein full name:

glyceraldehyde-3-phosphate dehydrogenase

# Synonyms:

1H18

## Immunogen:

Recombinant protein

# Isotype:

IgG

# Subcellular location:

Cytoplasm

#### **Purity:**

Affinity purification

### Form:

Liquid

## Storage Buffer:

PBS with 0.02%sodium azide,100  $\mu$ g/ml BSA and 50% glyce rol.


## Storage:

Store at -20 °C for one year.

# Experimental procedure:

Antigen retrieval: Citrate buffer (pH 6.0), Medium high heat for 8 minutes, stop for 7 minutes, medium high heat for 8 minutes. Incubate antibody, 4°C overnight. Secondary antibody: P oly-HRP Goat Anti-Rabbit & Mouse Universal Secondary Antibody, RT, 1h.

# Images:



Dilution of 1:50000 incubated at room temperature for 1.5 ho urs.

# Source of Reagents:

发表[中文论文]请标注:GAPDH(B31001T)由杭州浩克生物技术有限公司提供;



发表[英文论文]请标注:GAPDH(B31001T) wer e kindly provided by Hangzhou Haoke Biotechno logy Co., Ltd.